International Journal of Bifurcation and Chaos (C) World Scientific Publishing Company

# Imaginary Scators Bound Set Under The Iterated Quadratic Mapping In 1+2D Dynamical Space

M. FERNÁNDEZ-GUASTI<sup>\*</sup>

Lab. de Óptica Cuántica, Depto. de Física, Universidad A. Metropolitana - Iztapalapa, 09340 México D.F., Ap. postal. 55-534, MEXICO mfg@xanum.uam.mx, url: http://luz.izt.uam.mx

Received (to be inserted by publisher)

The quadratic iteration is mapped within imaginary scator algebra in 1+2 dimensions in dynamic space. There is a bound K set in the scator three dimensional space with a highly complex boundary  $J = \partial K$ . Two and three dimensional renderings of the K set exhibit a rich fractal boundary in all three directions. The Julia and filled in Julia sets are identically reproduced at two perpendicular planes where only one non-vanishing hypercomplex director component is present. The fixed points of K in  $\mathbb{S}^{1+2}$  can be obtained from the roots of a quadratic equation. In  $\mathbb{S}^{1+2}$  there can be hypercomplex roots that give rise to four additional fixed points. The K set of the origin is equal to the unit magnitude scator surface, in analogy to the unit circle Julia set of the origin. The ix set, is the three dimensional equivalent of the M-set in three dimensions. It is conjectured that the ix-set with some restrictions, is the set of parameters where the J set is connected.

*Keywords*: Hyper-complex numbers; Imaginary scators; Quadratic iteration; Julia set; Discrete dynamical systems.

# 1. Introduction

The iterated quadratic mapping in the cuspheric scator set, produces a rich and complex structure in parameter three dimensional space [Fernández-Guasti, 2016]. In contrast, other algebraic structures with dimension higher than two, such as quaternions, produce a disappointing surface of revolution of the M-set in three dimensions [Gomatam *et al.*, 1995; Bedding & Briggs, 1995]. There have other efforts to extend two dimensional fractal structures to three dimensions such as ternary algebra [Cheng & Tan, 2007] and triplex algebra [White & Nylander, 2009; Rama & Mishra, 2011] and variations of them [Bonzini, 2010]. Visualizations of these sets, in particular quasi-Fuschian fractals [Araki, 2006] and the mandelbulb, have received wide public exposure [Aron, 2009; Sanderson, 2009].

Methods originated in geographic visualization, architecture and digital animation are proving quite useful to render these rather complicated structures [Blackledge, 2002].

<sup>\*</sup>permanent address of the author.

In this communication, the three dimensional product and addition operations of imaginary scator algebra are invoked to evaluate the quadratic mapping in dynamic space. Scator algebra is a finite dimensional algebra over the reals with a multiplicative identity, thus fulfilling the hypercomplex algebra Kantor and Solodovnikov criteria except for the distributivity of the product over addition [Kantor & Solodovnikov, 1989]. Scator elements can thus be viewed as hypercomplex numbers in 1+n dimensions if the distributivity condition is relaxed. For n = 2, many of the singular properties of this algebra are already present. The scator product is commutative and all elements except zero and infinity have an inverse. Nonetheless, scator algebra is no longer a division algebra because it has zero products of non zero factors. The scator product is endowed with a second order involution. This feature can be used to establish an order parameter. These peculiarities do not prevent the scator number system from generating consistent iterated quadratic mappings with a rich fractal structure in parameter space [Fernández-Guasti, 2016].

Scator algebra also produces remarkable iterated quadratic mappings in dynamic space. This is the subject matter of the present communication: In section 2, the essentials of imaginary scator algebra in 1+2 dimensions are presented. The scator quadratic mapping is described in section 3. Three dimensional analogues of the Julia and filled in Julia sets are presented in this section. 2D and 3D renderings illustrate some of the main features of these sets. Square nilpotent points are discussed in subsection 3.2. The fingerprint of the origin is shown in subsection 3.3. Fixed points in scator dynamic space are described in section 4; hypercomplex roots produce extra fixed points with very interesting properties that coexist with the real or imaginary fixed points. Inverse orbits are discussed in section 5, the existence of four possible roots permits the evaluation of many points in the boundary set. Self similar objects within the K set are set forth in section 6. An example is expounded in section 7, where the K set exhibits a Cantor dust structure. Section 8 is dedicated some conjectures, given the difficulties encountered to establish formal proofs. Conclusions are drawn in the last section.

### 2. Imaginary scators

In the additive representation, scator elements in 1+2 dimensions are written as the sum of three components,

$$\overset{o}{\varphi} = s + x \,\check{\mathbf{e}}_x + y \,\check{\mathbf{e}}_y,\tag{1}$$

where  $s, x, y \in \mathbb{R}$  and  $\check{\mathbf{e}}_x, \check{\mathbf{e}}_y \notin \mathbb{R}$ . The first component is the *scalar component*, while subsequent components are named *director components* [Fernández-Guasti & Zaldívar, 2013]. Scator elements are decorated with an oval placed overhead<sup>1</sup>. Addition of scators is defined by the sum of coefficients in each component  $\overset{o}{\alpha} + \overset{o}{\beta} = (a_0 + a_x \check{\mathbf{e}}_x + a_y \check{\mathbf{e}}_y) + (b_0 + b_x \check{\mathbf{e}}_x + b_y \check{\mathbf{e}}_y) = (a_0 + b_0) + (a_x + b_x), \check{\mathbf{e}}_x + (a_y + b_y) \check{\mathbf{e}}_y$ . Scator numbers satisfy commutative group properties under addition in  $\mathbb{R}^{1+2}$ .

**Definition 2.1.** The extended scator product  $\overset{o}{\alpha}\overset{o}{\beta}$  of two scators,  $\overset{o}{\alpha} = a_0 + a_x \, \check{\mathbf{e}}_x + a_y \, \check{\mathbf{e}}_y$  and  $\overset{o}{\beta} = b_0 + b_x \, \check{\mathbf{e}}_x + b_y \, \check{\mathbf{e}}_y$  is,

$$\overset{o}{\alpha} \overset{o}{\beta} = a_0 b_0 \left( 1 - \frac{a_x b_x}{a_0 b_0} \right) \left( 1 - \frac{a_y b_y}{a_0 b_0} \right) + a_0 b_0 \left( 1 - \frac{a_y b_y}{a_0 b_0} \right) \left( \frac{a_x}{a_0} + \frac{b_x}{b_0} \right) \check{\mathbf{e}}_x + a_0 b_0 \left( 1 - \frac{a_x b_x}{a_0 b_0} \right) \left( \frac{a_y}{a_0} + \frac{b_y}{b_0} \right) \check{\mathbf{e}}_y.$$
(2)

If there are coefficients equal to zero in the scator factors, the director coefficients limits should be taken prior to the scalar coefficients limit.

 $<sup>^{1}</sup>$  \overset{o} in LATEX lore

The scator product is usually defined in the  $S^{1+2}$  set, where infinity is avoided. The extended scator product allows for divergent products.

**Example 2.1.** If  $a_x = b_x = 0$ ,

$$\lim_{a_x\to 0, b_x\to 0,} \left(\stackrel{o}{\alpha}\stackrel{o}{\beta}\right) = (a_0b_0 - a_yb_y) + (a_yb_0 + a_0b_y)\check{\mathbf{e}}_y.$$

The usual complex product in  $\mathbb{C}$  is recovered. The imaginary unit is  $\check{\mathbf{e}}_y$ . An analogous result is obtained if  $a_y = b_y = 0$ , but the imaginary unit is then  $\check{\mathbf{e}}_x$ .

**Example 2.2.** The limit  $a_0 \to 0$  if  $a_x a_y \neq 0$ , gives divergent coefficients in all three components,

$$\lim_{a_0\to 0} \begin{pmatrix} o & o \\ \alpha & \beta \end{pmatrix} = \infty + \infty \check{\mathbf{e}}_x + \infty \check{\mathbf{e}}_y.$$

There are then scator factors with finite coefficients whose product admits infinite coefficients.

**Example 2.3.** If  $a_0 = 0$  and  $a_x = 0$ . The director coefficient limit should be taken first, thus

$$\lim_{a_x \to 0} \left( \overset{o}{\alpha} \overset{o}{\beta} \right) = (a_0 b_0 - a_y b_y) + (a_0 b_0 - a_y b_y) \left( \frac{b_x}{b_0} \right) \check{\mathbf{e}}_x + (a_y b_0 + a_0 b_y) \check{\mathbf{e}}_y;$$

Thereafter, the scalar component limit is evaluated

$$\lim_{a_0 \to 0} \left( \lim_{a_x \to 0} \begin{pmatrix} o & o \\ \alpha & \beta \end{pmatrix} \right) = -a_y b_y - a_y b_y \frac{b_x}{b_0} \check{\mathbf{e}}_x + a_y b_0 \check{\mathbf{e}}_y.$$
(3)

**Example 2.4.** If one factor has vanishing director coefficients  $\overset{o}{\alpha} = a_0 + 0 \, \check{\mathbf{e}}_x + 0 \, \check{\mathbf{e}}_y = a_0$ ,

$$\overset{o}{\alpha}\overset{o}{\beta} = a_0\overset{o}{\beta} = a_0b_0 + a_0b_x\check{\mathbf{e}}_x + a_0b_y\check{\mathbf{e}}_y.$$

The  $a_0$  component produces a scaling of all the scator  $\stackrel{o}{\beta}$  components. Hence it is rightly named, the scalar component of the scator.  $\stackrel{o}{1} = 1 + 0 \, \check{\mathbf{e}}_x + 0 \, \check{\mathbf{e}}_y = 1$  is the multiplicative neutral.

**Example 2.5.** If  $a_x b_x = a_y b_y = a_0 b_0$ , then  $\overset{o}{\alpha} \overset{o}{\beta} = 0 + 0 \check{\mathbf{e}}_x + 0 \check{\mathbf{e}}_y$ , the scator product is zero. Thus, there exist zero products of non zero factors.

# 2.1. Multiplicative representation of scators

The multiplicative representation of scators is analogous to the polar representation of complex numbers in 1+1 dimensions,

$$\overset{o}{\varphi} = \varphi_0 e^{\varphi_x \, \check{\mathbf{e}}_x} e^{\varphi_y \, \check{\mathbf{e}}_y},\tag{4}$$

where e is the complex exponential function.  $\varphi_0$  is the scator magnitude also named the multiplicative scalar,  $\varphi_x$  is the angle of the scator projection between s and the  $\check{\mathbf{e}}_x$  axes,  $\varphi_y$  is the angle of the projection between s and the  $\check{\mathbf{e}}_y$  axes. In the multiplicative representation, the product of two scators is evaluated by performing the product of the magnitudes and the addition of the multiplicative director coefficients with the same director unit [Fernández-Guasti, 2021a],

$$\overset{o}{\alpha}\overset{o}{\beta} = \left(\alpha_0 e^{\alpha_x \check{\mathbf{e}}_x} e^{\alpha_y \check{\mathbf{e}}_y}\right) \left(\beta_0 e^{\beta_x \check{\mathbf{e}}_x} e^{\beta_y \check{\mathbf{e}}_y}\right) = \alpha_0 \beta_0 e^{(\alpha_x + \beta_x) \check{\mathbf{e}}_x} e^{(\alpha_y + \beta_y) \check{\mathbf{e}}_y}.$$
(5)

The multiplicative scator components having the same director unit  $\check{\mathbf{e}}_x$  or  $\check{\mathbf{e}}_y$ , satisfy the addition theorem for exponents. However, the addition theorem for exponents does not hold for scators with different director units, i.e.  $e^{\varphi_x \check{\mathbf{e}}_x} e^{\varphi_y \check{\mathbf{e}}_y} \neq e^{\varphi_x \check{\mathbf{e}}_x + \varphi_y \check{\mathbf{e}}_y}$ .

The additive (1) and multiplicative (4) representations of scators are related by

$$\overset{o}{\varphi} = \varphi_0 e^{\varphi_x \, \check{\mathbf{e}}_x} e^{\varphi_y \, \check{\mathbf{e}}_y} = \varphi_0 \cos \varphi_x \cos \varphi_y + \varphi_0 \cos \varphi_y \sin \varphi_x \, \check{\mathbf{e}}_x + \varphi_0 \cos \varphi_x \sin \varphi_y \, \check{\mathbf{e}}_y = s + x \, \check{\mathbf{e}}_x + y \, \check{\mathbf{e}}_y. \tag{6}$$

From this expression, it follows that if  $s = \varphi_0 \cos \varphi_x \cos \varphi_y = 0$ , then  $\overset{o}{\varphi} = 0$  or  $\overset{o}{\varphi}$  has only one non vanishing director component. Furthermore, if the two director components do not vanish,  $\varphi_0 \cos \varphi_y \sin \varphi_x \neq 0$  and  $\varphi_0 \cos \varphi_x \sin \varphi_y \neq 0$ , then the scalar component is necessarily also different from zero.

**Definition 2.2.** The scator set  $\mathbb{S}^{1+2} \subset \mathbb{R}^3$ , is the subspace where the scalar component is not zero if the two director components are different from zero,

$$\mathbb{S}^{1+2} = \left\{ \stackrel{o}{\varphi} = s + x \,\check{\mathbf{e}}_x + y \,\check{\mathbf{e}}_y, \ s, x, y \in \mathbb{R} : s \neq 0 \text{ if } x, y \neq 0 \right\}.$$
(7)

The  $\mathbb{S}^{1+2}$  set avoids divergent products (like the one in Example 2.2). This set was labeled with the letter  $\mathbb{E}$  in some of the earlier manuscripts.

Remark 2.1. The scator additive and multiplicative representations are equivalent in the  $\mathbb{S}^{1+2}$  scator set, except for the kernel of the transformation.

# 2.2. Magnitude

The conjugate of the scator is given by the negative of the director components, leaving the scalar component unchanged in either representation. In the additive representation,  $\overset{o^*}{\varphi} = s - x\check{\mathbf{e}}_x - y\check{\mathbf{e}}_y$  is the conjugate of  $\overset{o}{\varphi} = s + x\check{\mathbf{e}}_x + y\check{\mathbf{e}}_y$ . In the multiplicative representation,  $\overset{o^*}{\varphi} = \varphi_0 e^{-\varphi_x}\check{\mathbf{e}}_x e^{-\varphi_y}\check{\mathbf{e}}_y$  is the conjugate of  $\overset{o}{\varphi} = \varphi_0 e^{\varphi_x}\check{\mathbf{e}}_x e^{\varphi_y}\check{\mathbf{e}}_y$ . The magnitude of a scator is equal to the positive square root of the scator times its conjugate  $\|\overset{o}{\varphi}\| = \sqrt{\overset{o}{\varphi}} \overset{o^*}{\varphi}$ . In the multiplicative representation from (5), is  $\|\overset{o}{\varphi}\| = \varphi_0$ , the multiplicative scalar is thus the scator magnitude. In the additive representation, the square magnitude is

$$\left\| \overset{o}{\varphi} \right\|^2 = \overset{o}{\varphi} \overset{o}{\varphi}^* = s^2 \left( 1 + \frac{x^2}{s^2} \right) \left( 1 + \frac{y^2}{s^2} \right). \tag{8}$$

The multiplicative inverse, from (8) is  $\overset{o}{\varphi}^{-1} = \overset{o}{\varphi}^* \| \overset{o}{\varphi} \|^{-2}$ . In multiplicative variables  $\overset{o}{\varphi}^{-1} = \overset{o}{\varphi}^* / \varphi_0^2$  and in terms of additive variables, the inverse is

$$\overset{o}{\varphi}^{-1} = s^{-2} \left( 1 + \frac{x^2}{s^2} \right)^{-1} \left( 1 + \frac{y^2}{s^2} \right)^{-1} \overset{o}{\varphi}^*.$$
(9)

A scator is invertible except if all additive components are zero or if s = 0 when  $xy \neq 0$ .

**Definition 2.3.** The extended scator set  $\bar{\mathbb{S}}^{1+2} = \mathbb{R}^3 \cup \{\infty\}$ , allows for all three scator coefficients to be in the reals plus infinity

$$\bar{\mathbb{S}}^{1+2} = \left\{ \stackrel{o}{\varphi} = s + x \,\check{\mathbf{e}}_x + y \,\check{\mathbf{e}}_y, \ s, x, y \in \mathbb{R} \cup \{\infty\} \right\},\tag{10}$$

The points at infinity are attained if any of the scator components becomes infinite, i.e.  $s \to \infty$  and/or  $x \to \infty$  and/or  $y \to \infty$ . The magnitude of these points is infinite. Although the set of points with zero scalar component s = 0 and two nonvanishing director components  $xy \neq 0$  have finite coefficients, their magnitude is also infinite. Elements in  $\bar{S}^{1+2} \setminus S^{1+2}$  have divergent magnitude. This state of affairs has led to the concept of *divergent vicinity* whenever the zero scator is involved [Fernández-Guasti, 2016]. The element  $0 + 0 \check{\mathbf{e}}_x + 0 \check{\mathbf{e}}_y$  has a divergent vicinity since the magnitude of  $0 + \delta x \check{\mathbf{e}}_x + \delta y \check{\mathbf{e}}_y$  is infinite if  $\delta x, \delta y \neq 0$ . The scator magnitude does not diverge in  $S^{1+2}$ .

Remark 2.2. There exists a scator multiplicative representation for elements in the  $\mathbb{S}^{1+2}$  scator set but not in  $\overline{\mathbb{S}}^{1+2} \setminus \mathbb{S}^{1+2}$ .

Let us mention two propositions that will be useful in the coming sections:

**Lemma 1** [Fernandez-Guasti, 2016]. The only non trivial square nilpotent elements  $\overset{o^2}{\varphi} = 0 + 0 \,\check{\mathbf{e}}_x + 0 \,\check{\mathbf{e}}_y$  in 1+2 dimensional imaginary scator algebra are elements whose three components have equal absolute value,  $\overset{o}{\varphi} = \pm g \pm g \,\check{\mathbf{e}}_x \pm g \,\check{\mathbf{e}}_y$ .

**Corollary 2.1** [Fernandez-Guasti, 2016]. The square of an invertible element is invertible if it is not square nilpotent.

# 3. Iterated quadratic mapping

Consider a function  $\stackrel{o}{f_c}: \bar{\mathbb{S}}^{1+2} \to \bar{\mathbb{S}}^{1+2}$  to be a scator function of scator variable.

**Definition 3.1.** The non divergent set  $K(f_c)$  in dynamic space for imaginary scators in 1+2 dimensions is given by

$$K(\stackrel{o}{f}_{c}) = \left\{ \stackrel{o}{\varphi} \in \bar{\mathbb{S}}^{1+2} : \forall m \in \mathbb{N}, \left\| \stackrel{o}{f}_{c} \stackrel{o}{\varphi} \right\| \nrightarrow \infty \right\},$$
(11)

where  $\overset{o^m}{f_c}$  denotes the *m*-fold composition  $\overset{o^m}{f_c} = \overset{o}{f_c} (\overset{o}{f_c} (\overset{o}{f_c} \cdots))$  of the function  $\overset{o}{f_c} \in \mathbb{S}^{1+2}$ .

 $K(\mathring{f}_c)$  is a three dimensional generalization of the filled in Julia set. In this generalization, the Julia set is again defined as the boundary of the filled in Julia set  $J(\mathring{f}_c) = \partial K(\mathring{f}_c)$ . This boundary is now embedded in an  $\mathbb{S}^{1+2}$  three dimensional space. The filled in  $K(\mathring{f}_c)$  set has been defined in terms of a non divergent condition rather than a bounded one. This will be relevant as we shall see further down because nilpotent points can be very large but are nonetheless in the K set.

The square of a scator  $\overset{o}{\varphi} = s + x\check{\mathbf{e}}_x + y\check{\mathbf{e}}_y$  is obtained from the product of two equal scators,  $\overset{o}{\varphi}^2 = (s + x\check{\mathbf{e}}_x + y\check{\mathbf{e}}_y)^2 = s_\diamond + x_\diamond\check{\mathbf{e}}_x + y_\diamond\check{\mathbf{e}}_y$ . From Eq. (2)

$$\overset{o}{\varphi}^{2} = s^{2} \left( 1 - \frac{x^{2}}{s^{2}} \right) \left( 1 - \frac{y^{2}}{s^{2}} \right) + 2sx \left( 1 - \frac{y^{2}}{s^{2}} \right) \check{\mathbf{e}}_{x} + 2sy \left( 1 - \frac{x^{2}}{s^{2}} \right) \check{\mathbf{e}}_{y}.$$
 (12)

The square function mapping  $q: \mathbb{S}^{1+n} \to \mathbb{S}^{1+n}, \overset{o}{\varphi} \mapsto \overset{o^2}{\varphi}$  satisfies  $q(\lambda \overset{o}{\varphi}) = \lambda^2 q(\overset{o}{\varphi}), \lambda \in \mathbb{R}$ . In the multiplicative representation, the square of a scator is

$$\overset{o}{\varphi}^{2} = \varphi_{0}^{2} e^{2\varphi_{x}\check{\mathbf{e}}_{x}} e^{2\varphi_{y}\check{\mathbf{e}}_{y}}.$$
(13)

The iteration procedure in dynamic space is obtained by fixing an initial constant  $\overset{o}{c} = s_c + x_c \check{\mathbf{e}}_x + y_c \check{\mathbf{e}}_y$ and evaluating, for each point in the scator space  $\overset{o}{\varphi} = s + x \check{\mathbf{e}}_x + y \check{\mathbf{e}}_y \in \mathbb{S}^{1+2}$ , the quadratic recurrence relationship

$$\overset{o}{\varphi}_{m+1} = \overset{o}{\varphi}_m^2 + \overset{o}{c}. \tag{14}$$

Bound points obtained for an arbitrary number of iterations comprise the filled in Julia set in  $\bar{\mathbb{S}}^{1+2}$ . Consider the family of maps  $\stackrel{o}{f}_c: \stackrel{o}{\varphi} \mapsto \stackrel{o}{\varphi}^2 + \stackrel{o}{c}$  from  $\bar{\mathbb{S}}^{1+2}$  to  $\bar{\mathbb{S}}^{1+2}$ , where the variable  $\stackrel{o}{\varphi}$  and the constant  $\stackrel{o}{c}$  are scator elements. In terms of the additive coefficients, for each point  $\stackrel{o}{\varphi}_1 = s_1 + x_1 \check{\mathbf{e}}_x + y_1 \check{\mathbf{e}}_y$  and  $\stackrel{o}{c} = c_s + c_x \check{\mathbf{e}}_x + c_y \check{\mathbf{e}}_y$ , the quadratic iteration recurrence relationship for the scalar component is

$$s_{m+1} = s_m^2 \left( 1 - \frac{x_m^2}{s_m^2} \right) \left( 1 - \frac{y_m^2}{s_m^2} \right) + c_s$$
(15a)

and for the director components,

$$x_{m+1} = 2s_m x_m \left(1 - \frac{y_m^2}{s_m^2}\right) + c_x,$$
 (15b)

$$y_{m+1} = 2s_m y_m \left( 1 - \frac{x_m^2}{s_m^2} \right) + c_y.$$
 (15c)

The iteration (15a)-(15c) has been implemented in the Mandelbulber (version 2.20-dev) three dimensional fractal visualization program. Very intricate surfaces are obtained that are difficult to handle by ray tracing programs. The processing power required to generate 3D fractals is greatly increased compared with 2D renderings. It is necessary to extend the two dimensional image grid to three dimensional space. An image with  $10^3$  pixels resolution in each axis, requires  $10^9$  voxels (1000 million points!). Thus, the iteration of a much larger set of points is required to begin with. Thereafter, ray tracing is necessary to establish the observer point of view, with the concomitant occlusion of interior points or points that lie behind a particular surface from the observer point of view. The capability of using parallel GPU processing using OpenCL has greatly enhanced the rendering velocity.

Two dimensional renderings of the 3D  $K(\tilde{f}_c)$  set can also be made at a particular 2D plane with standard plots. For example, the set in the  $(s, \check{\mathbf{e}}_y)$  plane at a constant  $x_0\check{\mathbf{e}}_x$  value is obtained by evaluating the (15a)-(15c) triad for  $s + x_0\check{\mathbf{e}}_x + y\check{\mathbf{e}}_y$ , for all s, y values. Inclined planes can also be rendered if a condition of the form x = my, where m is constant, is imposed. Notice that in these cases, the plane under consideration has no breath, i.e. it is fixed to a single value. In contrast, 3D renderings require voxels, that is 3D tiny boxes; in this case, a plane can thus be obtained only within the thickness of the minimum voxel size.

# 3.1. Non divergent set in dynamic space for $\overset{o}{\varphi}_{c} = -0.5 + 0 \, \check{\mathrm{e}}_{x} + 0.5 \, \check{\mathrm{e}}_{y}$

Consider the quadratic iteration non divergent set in dynamic  $\mathbb{S}^{1+2}$  scator space  $K(\overset{o}{\varphi}^2 + \overset{o}{c})$  for the hypercomplex point  $\overset{o}{c} = -0.5 + 0 \,\check{\mathbf{e}}_x + 0.5 \,\check{\mathbf{e}}_y$ . The rendered set, observed from  $\overset{o}{v} = 3 - 8 \,\check{\mathbf{e}}_x + 1 \,\check{\mathbf{e}}_y$ , is shown in figure 1. This set is equivalent to the filled in Julia set in the complex plane  $K(z^2 + z_c) \in \mathbb{C}, z_c = -0.5 + 0.5i$  but



Fig. 1: Non divergent set in 3D dynamic scator space,  $K(\overset{o}{\varphi}^2 + \overset{o}{\varphi}_c) \in \mathbb{S}^{1+2}$  of the  $\overset{o}{c} = -0.5 + 0 \,\check{\mathbf{e}}_x + 0.5 \,\check{\mathbf{e}}_y$  hypercomplex point, seen from  $\overset{o}{v} = 3 - 8 \,\check{\mathbf{e}}_x + 1 \,\check{\mathbf{e}}_y$ . Image generated with 'Mandelbulber' rendering program.

extended to three dimensional space. The fractal nature of the surface makes it very difficult to produce an accurate ray tracing reflection due to the boundary roughness. Besides the central bundle, there are eight straight long filaments extending from the center up to large coordinate values. This feature can be explained as follows.

### 3.2. Square nilpotent elements

Square nilpotent  $\overset{o}{\varphi}_{nil}$  points are equal to the constant  $\overset{o}{c}$  upon the first iteration,

$$\dot{\varphi}_2 = \dot{\varphi}_{nil}^2 + \ddot{c} = \ddot{c}.$$

Thereafter, the iterations follow the orbit of the point  $\overset{o}{c}$  in parameter space, i.e.  $\{\overset{o}{\varphi}_{nil}, \overset{o}{c}, \overset{o}{c}^2 + \overset{o}{c}, (\overset{o}{c}^2 + c)^2 + \overset{o}{c}, ...\}$ . This sequence does not diverge if the point belongs to the Ix-set. The orbit of all nilpotent points after the first iteration is the same and is the sequence of  $\overset{o}{c}$  in parameter space. This set, coined here as the *Ix-set*<sup>2</sup>, has been described in an earlier communication [Fernández-Guasti, 2016]. The *Ix-set* is the quadratic iteration bound set in parameter  $\mathbb{S}^{1+2}$  scator space. The Mandelbrot set is a subset of the Ix-set,  $M \subset Ix$ . If the constant  $\overset{o}{c}$  is in the Ix-set, it is a bound point and therefore nilpotent points will be in the  $K(\overset{o}{\varphi}^2 + \overset{o}{c})$  set. From Lemma 1, nilpotent elements have equal absolute value of the scalar and director components. The four  $\overset{o}{\varphi} = \pm g \pm g \check{\mathbf{e}}_x \pm g \check{\mathbf{e}}_y$  straight lines are tilted  $\pm 45^\circ$  with respect to the *s*,  $\check{\mathbf{e}}_x$  and *s*,  $\check{\mathbf{e}}_y$  projections. The azimuthal angle with respect to any of the axes is  $\arctan(\pm\sqrt{2}) \approx \pm 54.7^\circ$ . The four spikes present in the  $K(\overset{o}{\varphi}^2 + \overset{o}{c})$  sets for  $\overset{o}{c} \in Ix$ -set, are due to this nilpotent feature. These spikes extend to arbitrarily large values of *g*.



(a) Rendering for values within  $\pm 10$  in all three variables.



(b) Rendering restricted to the  $0 \le x \le 10$  region in the  $\check{\mathbf{e}}_x$  axis.

Fig. 2:  $K(\overset{o}{\varphi}^2 + \overset{o}{c})$  closer detail of the  $\overset{o}{c} = -0.5 + 0 \, \check{\mathbf{e}}_x + 0.5 \, \check{\mathbf{e}}_y$  hypercomplex point, seen from  $\overset{o}{v} = 0.75 - 2 \, \check{\mathbf{e}}_x + 1 \, \check{\mathbf{e}}_y$ .

A close up of the central region is shown in figure 2a. The surface exhibits an intricate pattern of streaks in different directions. There is, so far, hardly any resemblance of this figure with a filled in Julia set. It is possible to limit the rendering to a region in the mandelbulber program. In figure 2b, the set is limited to  $0 \le x \le 10$  region in the  $\check{\mathbf{e}}_x$  direction and left in the default  $\pm 10$  limits in the scalar and  $\check{\mathbf{e}}_y$  axes. The set inside colouring, established by the programmers, is determined by the value at the end of the orbit trap iterations. The surface cut at x = 0 begins to look a bit more familiar. If the observation point is located on the  $\check{\mathbf{e}}_x$  axis, figure 3a is obtained and if the rendering is further restricted to a region very close to the x = 0 plane, figure 3b is obtained. In a 3D rendering, the voxels have finite size in all three directions, in fact, they are usually cubic structures. This means that the x = 0 plane cannot be chosen with zero breath. It has to encompass a finite albeit small depth. Some kinks, particularly evident close to the s = 0 line, are a consequence of this finite depth. For comparison, a filled in Julia set  $K(z_c)$  for  $z_c = -0.5 + 0.5i$  in the complex plane is shown in figure 3c. The two sets are remarkably similar, the irrelevant inline colouring making for most of the difference. The minor differences, mainly observed close to s = 0, but present in the

<sup>&</sup>lt;sup>2</sup>Ix is pronounced ish, like in lavish.



(a)  $K(\varphi^{o^2} + c)$  seen from  $v = 0 - 2\check{\mathbf{e}}_x + 0\check{\mathbf{e}}_y$  restricted to  $0 \le x \le 10$ .



(c) Familiar filled in Julia set of the  $z_c = -0.5 + 0.5i$  point in the complex plane obtained from a standard 2D plot.



(b)  $K(\overset{o}{\varphi}^2 + \overset{o}{c})$  seen from  $\overset{o}{v} = 0 - 2\check{\mathbf{e}}_x + 0\check{\mathbf{e}}_y$  but restricted to the minimum voxel size in  $x, -0.003 \le x \le 0.003$ .



(d)  $K(\overset{o}{\varphi}^2 + \overset{o}{c})$  evaluated at the plane  $\overset{o}{\varphi} = s + 0.003\check{\mathbf{e}}_x + y\check{\mathbf{e}}_y$ , with a standard 2D plot.

Fig. 3:  $K(\varphi^{2} + c)$  detail of the  $c = -0.5 + 0 \,\check{\mathbf{e}}_{x} + 0.5 \,\check{\mathbf{e}}_{y}$  hypercomplex point. Comparison of a thin 3D slice seen from the  $\check{\mathbf{e}}_{x}$  direction, the Julia set in the complex plane and a 2D rendering with an offset of 0.003 in the  $\check{\mathbf{e}}_{x}$  hypercomplex plane.

whole frontier when looked in detail, can be attributed to the finite depth of the 3D rendering. To confirm this assertion, the voxel size is estimated to be  $6.1 \times 10^{-3}$  per side. The 3D program is then evaluating the x = 0 plane with  $\pm 3.05 \times 10^{-3}$  resolution. In figure 3d, a 2D rendering at the  $\overset{o}{\varphi} = x + 0.003 \,\check{\mathbf{e}}_x + y \,\check{\mathbf{e}}_y$ hypercomplex plane is depicted. This set exhibits the kinks absent in the Julia set, but present in the 3D rendering.

# 3.3. The fingerprint of the origin.

Consider the constant point to be the origin,  $\overset{o}{c} = 0 + 0 \check{\mathbf{e}}_x + 0 \check{\mathbf{e}}_y$ . The quadratic iteration is then

$$\overset{o}{\varphi}_{m+1} = \overset{o}{\varphi}_m^2$$

From the multiplicative to additive representations (6) and the square of a scator in the multiplicative representation (13), the square of a scator is

$$\varphi^{o^2} = \varphi_0^2 e^{2\varphi_x \,\check{\mathbf{e}}_x} e^{2\varphi_y \,\check{\mathbf{e}}_y} = \varphi_0^2 \left[ \cos\left(2\varphi_x\right) \cos\left(2\varphi_y\right) + \cos\left(2\varphi_y\right) \sin\left(2\varphi_x\right) \,\check{\mathbf{e}}_x + \cos\left(2\varphi_x\right) \sin\left(2\varphi_y\right) \,\check{\mathbf{e}}_y \right].$$

Its magnitude is  $\| \varphi^2 \| = \varphi_0^2$ . The doubling of angles produces a scator rotation (not an Euclidean rotation) in the  $s, \check{\mathbf{e}}_x$  and  $s, \check{\mathbf{e}}_y$  planes that does not alter the scator magnitude. A repeated iteration  $\overset{o}{\varphi}_{m+1} = \overset{o}{\varphi}_m^2$  will then make the magnitude increase indefinitely for  $\| \varphi^0 \| > 1$  and will decrease monotonically for  $\| \varphi^0 \| < 1$ .

The magnitude will be invariant for unit magnitude  $\| \overset{o}{\varphi} \| = 1$ . The set for the origin in dynamic space should then be the isometric surface. This is indeed the case, figure 4a shows the 3D rendering of the  $J(\overset{o}{\varphi}^2)$  set for  $\overset{o}{c} = 0 + 0 \, \check{\mathbf{e}}_x + 0 \, \check{\mathbf{e}}_y$ . For comparison, the cusphere surface, drawn for unit magnitude from Eq. (6), is shown in figure 4b. The  $J(\overset{o}{\varphi}^2)$  set is compared with the isometric surface rather than  $K(\overset{o}{\varphi}^2)$ because it is the boundary  $J = \partial K$  that is actually equal to the cusphere surface.



(a)  $J(\varphi^{o^2})$  seen from  $\overset{o}{v} = 1.4 - 2.5 \,\check{\mathbf{e}}_x + 0.7 \,\check{\mathbf{e}}_y$ .



(b) Cusphere drawn from parametric plot of Eq. (6) with  $\varphi_0 = 1$ 

Fig. 4: Bound set for the origin  $\overset{o}{c} = 0 + 0 \,\check{\mathbf{e}}_x + 0 \,\check{\mathbf{e}}_y$  and the cusphere rendered from the unit magnitude condition,  $\|\overset{o}{\varphi}\| = 1$ .

# 4. Fixed points

**Lemma 2.** The fixed points in  $\mathbb{S}^{1+2}$  scator space for the quadratic iteration  $\overset{\circ}{\varphi}_{m+1} = \overset{\circ}{\varphi}_m^2 + \overset{\circ}{c}$ , provided that the constant  $\overset{\circ}{c} = c + 0 \check{\mathbf{e}}_x + 0 \check{\mathbf{e}}_y$  is a scalar are: If |4c| > 1, (four hypercomplex roots)

$$\overset{o}{\varphi}_{\mathbb{S}^{1+2}\setminus\mathbb{S}^{1+1}} = \left(c + \frac{1}{4}\right) \pm \frac{1}{4}\sqrt{16c^2 - 1}\check{\mathbf{e}}_x \pm \frac{1}{4}\sqrt{16c^2 - 1}\check{\mathbf{e}}_y; \tag{16a}$$

If  $4c \leq 1$ , (real roots),

$$\overset{o}{\varphi}_{\mathbb{S}^{1+0}} = \frac{1}{2} \pm \frac{1}{2}\sqrt{1-4c};$$
 (16b)

If 4c > 1, (two copies of the complex like roots),

$$\overset{o}{\varphi}_{\mathbb{S}_x^{1+1}} = \frac{1}{2} \pm \frac{1}{2} \sqrt{4c - 1} \check{\mathbf{e}}_x, \quad \overset{o}{\varphi}_{\mathbb{S}_y^{1+1}} = \frac{1}{2} \pm \frac{1}{2} \sqrt{4c - 1} \check{\mathbf{e}}_y.$$
 (16c)

*Proof.* The point  $\overset{o}{\varphi}$  is fixed if  $\overset{o}{\varphi}_{m+1} = \overset{o}{\varphi}_m$  in the quadratic iteration. The polynomial to be solved is then  $\overset{o}{\varphi}^2 - \overset{o}{\varphi} + c = 0,$  (17)

where  $\overset{o}{\varphi} \in \mathbb{S}^{1+2}$ . The scator solutions to this polynomial are given by Theorem 1 in [Fernández-Guasti, 2021b]:

**Theorem 1** [Fernandez-Guasti, 2021]. The second order polynomial  $a \overset{o}{\varphi}^2 + b \overset{o}{\varphi} + c = 0$ , where  $\overset{o}{\varphi} \in \mathbb{S}^{1+2}$  is an elliptic scator and  $a, b, c \neq 0$  are real coefficients, has the following roots: If  $|4ac| > b^2$ ,

$$\overset{o}{\varphi}_{\mathbb{S}^{1+2}\setminus\mathbb{S}^{1+1}} = -\frac{4ac+b^2}{4ab} \pm \sqrt{\frac{(4ac)^2 - (b^2)^2}{16a^2b^2}} \check{\mathbf{e}}_x \pm \sqrt{\frac{(4ac)^2 - (b^2)^2}{16a^2b^2}} \check{\mathbf{e}}_y;$$
(18a)

If  $4ac \leq b^2$ ,

$$\overset{o}{\varphi}_{\mathbb{S}^{1+0}} = -\frac{b}{2a} \pm \frac{\sqrt{b^2 - 4ac}}{2a};$$
 (18b)

If  $4ac > b^2$ ,

$$\overset{o}{\varphi}_{\mathbb{S}_{1}^{1+1}} = -\frac{b}{2a} \pm \frac{\sqrt{-b^{2} + 4ac}}{2a} \check{\mathbf{e}}_{x}, \quad \overset{o}{\varphi}_{\mathbb{S}_{2}^{1+1}} = -\frac{b}{2a} \pm \frac{\sqrt{-b^{2} + 4ac}}{2a} \check{\mathbf{e}}_{y}.$$
(18c)

For the fixed points quadratic polynomial (17), a = 1, b = -1. From (18a)-(18c), the solutions (16a)-(16c) are obtained.

In contrast with the quadratic polynomial in the complex field, where the roots are either both real or both imaginary, the hypercomplex solutions can coexist with the real or complex like solutions in  $\mathbb{S}^{1+2}$ . We refer to hypercomplex roots to those solutions where the scalar and both director components are different from zero.

- If  $-\frac{1}{4} \le c \le \frac{1}{4}$ , only the usual two, possibly degenerate, real solutions exist. Outside this region, there are additionally four hypercomplex roots.
- For  $c < -\frac{1}{4}$ , there are six fixed points. In addition to the two real points, there are four fixed points due to the hypercomplex roots. These point lie in the  $\check{\mathbf{e}}_x, \check{\mathbf{e}}_y$  plane in the negative s semispace at  $s = c + \frac{1}{4}$ .
- For  $c > \frac{1}{4}$ , there are eight fixed points. Four fixed points, two in the  $s, \check{\mathbf{e}}_x$  plane and two in the  $s, \check{\mathbf{e}}_y$  plane. These points correspond to the usual solutions in the complex plane, but there are now two hypercomplex planes sharing the scalar axis. There is no precedence of the two hyperimaginary axes, the two hyperimaginary units  $\check{\mathbf{e}}_x$  or  $\check{\mathbf{e}}_y$  become identical to the *i* imaginary unit if only one (hyperimaginary) director component is present. In addition, there are four hypercomplex roots in the positive *s* semispace at the  $s = c + \frac{1}{4}$  plane. There exist hypercomplex roots ( $|4ac| > b^2$ ) whenever there exist complex like roots ( $4ac > b^2$ ).

The hypercomplex roots have director components with equal absolute value. Therefore, the four fixed points lie symmetrically placed in 45° planes with respect to the  $\check{\mathbf{e}}_x, \check{\mathbf{e}}_y$  axes. Just as in the complex plane, the constant c may be any point in  $c \in \mathbb{C}$ ; in  $\mathbb{S}^{1+2}$ , the constant  $\overset{\circ}{c}$  can be any point in the  $\mathbb{S}^{1+2}$  scator set. However, analytic solutions to the quadratic equation in  $\mathbb{S}^{1+2}$  are only available at present for scalar (real)  $\overset{\circ}{c}$ .

# 4.1. $c = -\frac{1}{2}$ , connected set

If  $\overset{o}{c} = -\frac{1}{2} + 0 \,\check{\mathbf{e}}_x + 0 \,\check{\mathbf{e}}_y$ , from (16b) the well known two real roots are obtained. These two fixed points  $\overset{o}{\varphi}_{\mathbb{S}^{1+0}} = \frac{1}{2} \pm \frac{\sqrt{3}}{2}$ , are shown in red in the complex plane 2D rendering in figure 5a. As usual, one of them is at the boundary  $\overset{o}{\varphi}_{\mathbb{S}^{1+0},+} = \frac{1}{2} + \frac{\sqrt{3}}{2} \in J(\overset{o}{\varphi}^2 - \frac{1}{2})$ , while the other is an interior point  $\overset{o}{\varphi}_{\mathbb{S}^{1+0},-} = \frac{1}{2} - \frac{\sqrt{3}}{2} \in K(\overset{o}{\varphi}^2 - \frac{1}{2}) \setminus J$ . This set is also shown in a 3D rendering in figure 5b. The  $K(\overset{o}{\varphi}^2 - \frac{1}{2})$  volume has been restricted to the positive semi-space, thus exhibiting the x = 0 plane in the forefront since the viewpoint is located on the  $\check{\mathbf{e}}_x$  axis at -2, i.e.  $\overset{o}{v}_{obs} = 0 - 2 \,\check{\mathbf{e}}_x + 0 \,\check{\mathbf{e}}_y$ .

Imaginary scators and quadratic mappings in three dimensions 11



(a) Julia set for  $c = -\frac{1}{2}$ . Fixed points in the real axis shown in red.



(b) 3D rendering of the  $K(\overset{o}{\varphi}^2 - \frac{1}{2})$  seen from  $\overset{o}{\nu}_{obs} = 0 - 2 \check{\mathbf{e}}_x + 0 \check{\mathbf{e}}_y$ , restricted to  $0 \le x \le 10$ 



(c) 2D rendering of the  $K\left(\varphi^{2}-\frac{1}{2}\right)$ , s = -0.25. The fixed points arising from hypercomplex roots shown in red.



(d) 3D rendering of the  $K(\overset{o^2}{\varphi} - \frac{1}{2})$  seen from  $\overset{o}{\varphi}_{obs} = 2 + 0 \,\check{\mathbf{e}}_x + 0 \,\check{\mathbf{e}}_y$ , restricted to  $-10 \leq x \leq -0.25$ 



(e)  $K(\varphi^2 - \frac{1}{2})$ , restricted to -10  $\leq s \leq -0.25$  and  $0 \leq x \leq 10$ .

Fig. 5: Non divergent set in  $\mathbb{S}^{1+2}$  dynamic space for  $c = -\frac{1}{2}$ .

In addition to these two real roots, the hypercomplex roots from (16a) are

$$\overset{o}{\varphi}_{\mathbb{S}^{1+2}\setminus\mathbb{S}^{1+1}} = -\frac{1}{4} \pm \frac{\sqrt{3}}{4}\check{\mathbf{e}}_x \pm \frac{\sqrt{3}}{4}\check{\mathbf{e}}_y.$$
(19)

These roots are depicted in red in figure 5c, where a two dimensional rendering of the  $K(\overset{o}{\varphi}^2 - \frac{1}{2})$  volume intersection with the  $\check{\mathbf{e}}_x, \check{\mathbf{e}}_y$  plane at s = -0.25 is shown depicted. These four hypercomplex roots, shown in red, belong to the J set since they lie on the boundary of the filled in set,  $J(\overset{o}{\varphi}^2 - \frac{1}{2}) = \partial K(\overset{o}{\varphi}^2 - \frac{1}{2})$ . This set is the higher dimensional analogue of the Julia set in the complex plane. In figure 5d, a 3D rendering is shown viewed from the s axis, the  $K(\overset{o}{\varphi}^2 - \frac{1}{2})$  set is limited to the interval  $-0.25 \ge s \ge -10$ . A perspective where two of the hypercomplex fixed points and one of the real fixed points are shown in red, is depicted in figure 5e. There are then 6 fixed points in  $K(\overset{o}{\varphi}^2 - \frac{1}{2})$  in  $\mathbb{S}^{1+2}$  scator space, in contrast with the 2 fixed points in  $\mathbb{S}^{1+1}$  scator space (equivalent to the complex plane).

### 5. Inverse orbits

The square roots of a scator  $\overset{o}{\varphi} \in \mathbb{S}^{1+2}$  in the additive representation with multiplicative angle variables are given by [Fernández-Guasti, 2022]

$$\overset{o}{\varphi}^{\frac{1}{2}} = \overset{o}{\zeta}_{\pm,0} = \pm \varphi_0^{\frac{1}{2}} \left( \cos \frac{\varphi_x}{2} \cos \frac{\varphi_y}{2} + \cos \frac{\varphi_y}{2} \sin \frac{\varphi_x}{2} \check{\mathbf{e}}_x + \cos \frac{\varphi_x}{2} \sin \frac{\varphi_y}{2} \check{\mathbf{e}}_y \right), \tag{20a}$$

and due to the  $\pi$ -pair symmetry,

$$\overset{o}{\varphi}^{\frac{1}{2}} = \overset{o}{\zeta}_{\pm,1} = \pm \varphi_0^{\frac{1}{2}} \left( \sin \frac{\varphi_x}{2} \sin \frac{\varphi_y}{2} - \sin \frac{\varphi_y}{2} \cos \frac{\varphi_x}{2} \check{\mathbf{e}}_x - \sin \frac{\varphi_x}{2} \cos \frac{\varphi_y}{2} \check{\mathbf{e}}_y \right).$$
(20b)

In terms of the scator additive variables  $\sqrt{\overset{o}{\varphi}} = \sqrt{s + x \,\check{\mathbf{e}}_x + y \,\check{\mathbf{e}}_y}$ , the square roots are

$$\sqrt{\overset{o}{\varphi}} = \overset{o}{\zeta}_{\pm,0} = \pm \frac{1}{2} \sqrt{\frac{1}{|s|}} \left[ \sqrt{\left(\sqrt{s^2 + x^2} + s\right) \left(\sqrt{s^2 + y^2} + s\right)} + \operatorname{Sgn} x \sqrt{\left(\sqrt{s^2 + x^2} - s\right) \left(\sqrt{s^2 + y^2} + s\right)} \, \check{\mathbf{e}}_x + \operatorname{Sgn} y \sqrt{\left(\sqrt{s^2 + x^2} + s\right) \left(\sqrt{s^2 + y^2} - s\right)} \, \check{\mathbf{e}}_y \right], \quad (21a)$$

and from the  $\pi$ -pair symmetry,

$$\sqrt{\frac{o}{\varphi}} = \overset{o}{\zeta}_{\pm,1} = \pm \frac{1}{2} \sqrt{\frac{1}{|s|}} \left[ \operatorname{Sgn} x \operatorname{Sgn} y \sqrt{\left(\sqrt{s^2 + x^2} - s\right) \left(\sqrt{s^2 + y^2} - s\right)} - \operatorname{Sgn} y \sqrt{\left(\sqrt{s^2 + x^2} + s\right) \left(\sqrt{s^2 + y^2} - s\right)} \check{\mathbf{e}}_x - \operatorname{Sgn} x \sqrt{\left(\sqrt{s^2 + x^2} - s\right) \left(\sqrt{s^2 + y^2} + s\right)} \check{\mathbf{e}}_y \right]. \quad (21b)$$

From Lemma 2, the hypercomplex roots of the quadratic equation (17) have equal director coefficients magnitudes,  $x^2 = y^2$  in (16a). Imposing this condition, the square roots (21a) are

$$\zeta_{\pm,0}^{o} = \pm \frac{1}{2} \sqrt{\frac{1}{|s|}} \left( \left( \sqrt{s^2 + x^2} + s \right) + x \, \check{\mathbf{e}}_x + y \, \check{\mathbf{e}}_y \right),$$
 (22a)

and (21b)

$$\overset{o}{\zeta_{\pm,1}} = \pm \frac{1}{2} \sqrt{\frac{1}{|s|}} \left( \operatorname{Sgn} x \operatorname{Sgn} y \left( \sqrt{s^2 + x^2} - s \right) - y \,\check{\mathbf{e}}_x - x \,\check{\mathbf{e}}_y \right), \tag{22b}$$

where x and y have been retained in the director coefficients to account for their sign. Invert equation (14) as usual to obtain the preimage of  $\overset{o}{\varphi}_{m+1}$ ,

$$\overset{o}{\varphi}_{m} = \sqrt{\overset{o}{\varphi}_{m+1} - \overset{o}{c}},$$

where the roots are given by (22a) and (22b).

# 5.1. Preimages of the $J(\varphi^{o^2} - \frac{1}{2})$ fixed points

The hypercomplex fixed points for  $\overset{o}{c} = -\frac{1}{2} + 0 \check{\mathbf{e}}_x + 0 \check{\mathbf{e}}_y$  are given by (19). Upon subtraction of  $\overset{o}{c}$ , the roots of

$$\overset{o}{\varphi}_{m}=\sqrt{rac{1}{4}\pmrac{\sqrt{3}}{4}}\check{\mathbf{e}}_{x}\pmrac{\sqrt{3}}{4}\check{\mathbf{e}}_{y},$$

need to be evaluated to find the inverse orbits of the fixed points. Consider one of the fixed points, say the one in the positive x and y quadrant,  $\overset{o}{\varphi}_{m+1} = \overset{o}{\varphi}_{inv1} = -\frac{1}{4} + \frac{\sqrt{3}}{4}\check{\mathbf{e}}_x + \frac{\sqrt{3}}{4}\check{\mathbf{e}}_y$ ,

$$\overset{o}{\varphi}' = s' + x' \check{\mathbf{e}}_x + y' \check{\mathbf{e}}_y = \overset{o}{\varphi}_{inv1} - \overset{o}{c} = \frac{1}{4} + \frac{\sqrt{3}}{4}\check{\mathbf{e}}_x + \frac{\sqrt{3}}{4}\check{\mathbf{e}}_y.$$

The square roots from (22a)-(22b) are then

$$\sqrt{\overset{o}{\varphi}_{inv1} - \overset{o}{c}} = \overset{o}{\zeta}_{\pm,0} = \pm \left(\frac{3}{4} + \frac{\sqrt{3}}{4}\,\check{\mathbf{e}}_x + \frac{\sqrt{3}}{4}\,\check{\mathbf{e}}_y\right),\,$$

and

$$\sqrt{\overset{o}{\varphi}_{inv1}} - \overset{o}{c} = \overset{o}{\zeta}_{\pm,1} = \pm \left(\frac{1}{4} - \frac{\sqrt{3}}{4}\check{\mathbf{e}}_x - \frac{\sqrt{3}}{4}\check{\mathbf{e}}_y\right).$$

From these four roots, the outcome of  $\overset{o}{\zeta}_{-,1}$  is again the fixed point  $\overset{o}{\varphi}_{inv1}$ ,

$$\sqrt{\overset{o}{\varphi}_{inv1} - \overset{o}{c}} = \left(\frac{1}{4} + \frac{\sqrt{3}}{4}\,\check{\mathbf{e}}_x + \frac{\sqrt{3}}{4}\,\check{\mathbf{e}}_y\right)^{\frac{1}{2}} = \overset{o}{\zeta}_{-,1} = -\frac{1}{4} + \frac{\sqrt{3}}{4}\,\check{\mathbf{e}}_x + \frac{\sqrt{3}}{4}\,\check{\mathbf{e}}_y = \overset{o}{\varphi}_{inv1}.$$

The fixed point  $\overset{o}{\varphi}_{inv1}$  is forward invariant (by definition) and backward invariant under the action of  $\overset{o}{\zeta}_{-,1}$ . The inverse orbit iteration evaluated with the  $\overset{o}{\zeta}_{+,0}$  root, seems to converge to a constant scalar component while the director components decrease monotonically, as shown in Table 1. A few iterates (red points)

$\zeta_{+,0}^{o}$	8	x	y
$\varphi_{inv1}^{o}$	-0.25	0.433013	0.433013
$\left( \begin{pmatrix} o \\ \varphi_{inv} - c \end{pmatrix}^{\frac{1}{2}} \right)$	0.75	0.433013	0.433013
÷	1.15062	0.193649	0.193649
	1.28917	0.0753636	0.0753636
	1.33819	0.0281712	0.0281712
	1.35588	0.0103891	0.0103891
	1.36232	0.00381307	0.00381307
		•	•
	$\frac{1}{2} + \frac{\sqrt{3}}{2} \approx 1.36603\ldots$	0	0

Table 1: Inverse orbit of fixed point  $\overset{o}{\varphi}_{inv1}$  evaluated with the  $\overset{o}{\zeta}_{+,0} = \frac{1}{2}\sqrt{\frac{1}{|s|}}\left(\sqrt{s^2 + x^2} + s + x\,\check{\mathbf{e}}_x + y\,\check{\mathbf{e}}_y\right)$  root. All points are in the J set.

are depicted in Figure 6, where the intersection of the  $K(\overset{o}{\varphi}^2 - \frac{1}{2})$  set with a plane having equal director components is pictured. The sequence seems to converge towards  $1.36...+0\check{\mathbf{e}}_x + 0\check{\mathbf{e}}_y$ , but interestingly,

#### $14 \quad M. \ Fern{and} ez-Guasti$

the value of the real fixed point in the Julia set is  $\overset{o}{\varphi}_{\mathbb{S}^{1+0},+} = \frac{1}{2} + \frac{\sqrt{3}}{2} \approx 1.36603\cdots$ . The point labeled 1.366 in figure 6 is the same point depicted in the boundary of Figures 5a and 5b. In these two figures, the plane intersecting  $K(\overset{o}{\varphi}^2 - \frac{1}{2})$  is x = 0 (or y = 0), whereas the intersecting plane is x = y in Figure 6.



Fig. 6: The  $K(\overset{o}{\varphi}^2 - \frac{1}{2})$  set observed at the intersection with the x = y plane (equal director components). The axes plot are s (abscissas) versus  $\frac{1}{\sqrt{2}}(\check{\mathbf{e}}_x + \check{\mathbf{e}}_y)$  (ordinates). Inverse orbits of the fixed point  $\overset{o}{\varphi}_{inv1} = -\frac{1}{4} + \frac{\sqrt{3}}{4}\check{\mathbf{e}}_x + \frac{\sqrt{3}}{4}\check{\mathbf{e}}_y$  are shown.

Backward iterations of  $\overset{o}{\varphi}_{inv1}$  with the  $\overset{o}{\zeta}_{+,1}$  root, give values with ever smaller scator coefficients in all three components as can be seen in Table 2. While the scalar component decreases monotonically, the director components alternate signs although their absolute value also decreases monotonically. These iterates are shown with cyan dots in Figure 6, the value of one iterate to the next joined by yellow lines. While the origin is seen as an interior point of the  $K(\overset{o}{\varphi}^2 - \frac{1}{2})$  set in the y = 0 plane, the indentations of this set in the x = y plane portray the origin as a boundary point.

Finally, backward iterations of  $\overset{o}{\varphi}_{inv1}$  with  $\overset{o}{\zeta}_{-,0} = -\frac{1}{2}\sqrt{\frac{1}{|s|}}\left(\left(\sqrt{s^2+x^2}+s\right)+x\,\check{\mathbf{e}}_x+y\,\check{\mathbf{e}}_y\right)$  produce a period two backward orbit alternating between the points

$$-\frac{1}{4} + \frac{\sqrt{3}}{4}\check{\mathbf{e}}_x + \frac{\sqrt{3}}{4}\check{\mathbf{e}}_y \longleftrightarrow -\frac{3}{4} - \frac{\sqrt{3}}{4}\check{\mathbf{e}}_x - \frac{\sqrt{3}}{4}\check{\mathbf{e}}_y$$

The  $\overset{o}{\varphi}_p = -\frac{3}{4} - \frac{\sqrt{3}}{4}\check{\mathbf{e}}_x - \frac{\sqrt{3}}{4}\check{\mathbf{e}}_y$  point is shown in magenta in Figure 6.

Since the J set is invariant under the iteration, points in the J set must map to points within the J set. We have not proved that the fixed hypercomplex points belong to the J set, although the fact that they are forward invariant but are neither zero nor infinity, suggest that they are in J. This conjecture is further supported by the visual position of these points in regions that seem to be on the boundary of the K set. All the points shown in Figure 6 again seem to lie on the  $\partial K = J$  boundary. Many other cases that are not reported also follow this pattern.

It is not mandatory to use the same root in the backward evaluation of orbits. We could proceed as in inverse Julia calculations in  $\mathbb{C}$ , where one of the two roots is randomly chosen. The random choice of one of the four hypercomplex roots in the present case, would fill in many other regions of the  $K(\overset{o}{\varphi}^2 - \frac{1}{2})$  set

$\zeta_{+,1}$	S	x	y
$\varphi_{inv1}$	-0.25	0.433013	0.433013
$\left( \begin{pmatrix} o \\ \varphi_{inv1} - c \end{pmatrix}^{\frac{1}{2}} \right)$	0.25	-0.433013	-0.433013
÷	0.0669873	0.25	0.25
	0.0349738	-0.166006	-0.166006
	0.0172025	0.113482	0.113482
	0.00855398	-0.0788981	-0.0788981
	0.00426558	0.0553182	0.0553182
	0.00213004	-0.0389501	-0.0389501
	0.00106434	0.0274834	0.0274834
			:

Table 2: Inverse orbit of fixed point  $\overset{o}{\varphi}_{inv1}$  evaluated with the  $\overset{o}{\zeta}_{+,1} = \frac{1}{2}\sqrt{\frac{1}{|s|}} \left( \operatorname{Sgn} x \operatorname{Sgn} y \left( \sqrt{s^2 + x^2} - s \right) - y \check{\mathbf{e}}_x - x \check{\mathbf{e}}_y \right)$  root. All points are in the J set, not interior points of the filled in Julia set,  $\notin K \setminus J$ .

intersection with the x = y plane (Figure 6). Notice that both, the forward or backward iterations maintain the equality between director coefficients, so all the iterated points coming from the hypercomplex roots must lie on this plane or its orthogonal version, x = -y.

Several interesting cases arise when applying different roots in the inverse orbit calculations. Let us undertake just one of them as a curious example. Consider the point  $\overset{o}{\varphi}_p = -\frac{3}{4} - \frac{\sqrt{3}}{4}\check{\mathbf{e}}_x - \frac{\sqrt{3}}{4}\check{\mathbf{e}}_y$  that is the partner of the fixed point  $\overset{o}{\varphi}_{inv1}$  in the period two backward orbit with the  $\overset{o}{\zeta}_{-,0}$  root. If the point  $\overset{o}{\varphi}_p$  is backward iterated with the  $\overset{o}{\zeta}_{-,1}$  root instead, the result is again  $\overset{o}{\varphi}_p$ . The point  $\overset{o}{\varphi}_p$  is then backwards invariant under  $\overset{o}{\zeta}_{-,1}!$  However it is not forward invariant, since the forward iteration maps it onto  $\overset{o}{\varphi}_{inv1}$ . Zero and infinity are attractors. Cannot prove



Fig. 7: Detail of the vicinity of 1.366 Increasing magnification of region around the real fixed point  $\frac{1}{2} + \frac{\sqrt{3}}{2} \approx 1.366$  observed at the plane with equal director components.

# 6. Self similarity

Self similarity at arbitrarily small scales is a hallmark of fractal structures. In the Ix-set ( $\mathbb{S}^{1+2}$  higher dimensional version of the M-set), we have seen that the main structure is repeated in very similar smaller structures. In dynamic space self-similarity in the complex plane often reproduces the form of the boundary or certain features at different scales rather than the whole object. Self similarity of the K set in  $\mathbb{S}^{1+2}$  in a region near to the scalar axis and one hyperimaginary plane is perhaps expected because such a plane is

close to the complex plane. For this reason, we have chosen to exhibit self similarity in a hyperimaginaryhyperimaginary plane region in Figure 8. The constant  $\stackrel{\circ}{c} = -1 + 0\check{\mathbf{e}}_x + 0\check{\mathbf{e}}_y$  has been set at -1. The part of the set shown in brown streaks is the surface of the volume between  $-10 \leq s < 0.6$ . The white-yelloworange cross like features correspond to constant s = 0.6, where the rendering of the set was stopped. The



(a) Viewpoint  $\overset{o}{v} = 2 + 0\check{\mathbf{e}}_x + 0\check{\mathbf{e}}_y$ .

(b) Detail of lower left branch.

(c) Further detail within branch.

Fig. 8: Self similarity of the  $K(\varphi^{o^2} - 1)$  set.  $\check{\mathbf{e}}_x, \check{\mathbf{e}}_y$  is seen from the *s* axis with  $s \leq 0.6$ . The cross shaped patterns on the s = 0.6 plane keep on repeating at different scales.

cross like motives are repeated along the diagonals with different sizes. The brown fibers intertwining is also repeated again and again but are more difficult to discern. Two close ups shown in Figure 8, keep on revealing more and more crossed motives at different scales.

\*\*\*\*\*\* The notation for the director coefficients  $x(m, r_x)$ ,  $y(m, r_y)$  follow a similar convention. The solutions have been labeled with different subindices in r because any of the  $m^3$  real roots combinations between them is possible.++

# 7. The $c = \frac{1}{2}$ , non connected set



(a)  $J\left(\dot{\varphi}^2 + \frac{1}{2}\right)$  set seen from  $2 - 4\check{\mathbf{e}}_x + 1\check{\mathbf{e}}_y$ 



(b)  $J\left(\dot{\varphi}^2 + \frac{1}{2}\right)$  set seen from  $0 - 4\check{\mathbf{e}}_x + 0\check{\mathbf{e}}_y$ 



(c)  $J\left(\overset{o^2}{\varphi} + \frac{1}{2}\right)$  set seen from  $0 - 4\check{\mathbf{e}}_x + 0\check{\mathbf{e}}_y$ , limited to  $0 \leq x \leq 0.03$ 



(d) Julia set for  $c = \frac{1}{2}$ .

Fig. 9: Visualization of the non divergent J set of the function  $\overset{o}{\varphi}^2 + c$ , with  $c = \frac{1}{2}$ .

Consider the non divergent set in dynamic space for the constant  $\stackrel{o}{c} = \frac{1}{2} + 0 \,\check{\mathbf{e}}_x + 0 \,\check{\mathbf{e}}_y$ . The 3D rendering, shown in figure 9a, reveals a grainy structure in contrast with the continuous rough surface of the previous

example. A Fatou dust like disconnected set in three dimensions is appreciated. Although there are some more or less dense regions, it is possible to 'see through' in any one of them. If  $c = \frac{1}{2}$ , recall that in  $\mathbb{C}$ , the Julia set is not connected since  $\frac{1}{2}$  is not in the M-set. The 3D rendering suggests that, in a similar fashion, since  $\frac{1}{2}$  is not in the Ix-set, the J set in  $\mathbb{S}^{1+2}$  is not connected. This remark will be discussed in more detail in section 8. Let us consider the well known  $z = \frac{1}{2} \pm \frac{1}{2}i$  two fixed points in  $\mathbb{C}$  to begin with; they are depicted in red in figure 9d. In  $\mathbb{S}^{1+2}$ , the corresponding two copies of these roots in the s,  $\check{\mathbf{e}}_x$  and  $s, \check{\mathbf{e}}_y$  planes are given by (16c),  $\overset{o}{\varphi}_{\mathbb{S}_x^{1+1}} = \frac{1}{2} \pm \frac{1}{2}\check{\mathbf{e}}_x$  and  $\overset{o}{\varphi}_{\mathbb{S}_y^{1+1}} = \frac{1}{2} \pm \frac{1}{2}\check{\mathbf{e}}_y$ . These last two roots are depicted in figure 9c. 3D renderings limited to thin regions are particularly difficult to visualize for non connected sets. In this example, if x is restricted to a 0.01 layer or less, there are hardly any points in the plot. As the thickness is increased, more bound points are present but the figure is increasingly less similar to a 2D rendering. A compromise has been chosen, where the layer has been restricted to  $0 \leq x \leq 0.03$ , in order to evince the similarities of the 3D and 2D mappings in figures 9c and the familiar 2D Julia set shown in figure 9d.

In addition to the two copies of the complex like roots, there are four hypercomplex roots. From (16a),

$$\overset{o}{\varphi}_{\mathbb{S}^{1+2}\setminus\mathbb{S}^{1+1}} = \frac{3}{4} \pm \frac{\sqrt{3}}{4}\check{\mathbf{e}}_x \pm \frac{\sqrt{3}}{4}\check{\mathbf{e}}_y.$$
(23)

These four fixed points are shown in red in the 2D plot in figure 10c. The 3D rendering of the  $K\left(\overset{o^2}{\varphi} + \frac{1}{2}\right)$ 



(a)  $J\left(\varphi^{o^2} + \frac{1}{2}\right)$  set seen from  $4 + 0\check{\mathbf{e}}_x + 0\check{\mathbf{e}}_y$ 



(b)  $J\left(\varphi^{o^2} + \frac{1}{2}\right)$  set restricted to  $0.74 \le s \le 0.75$ , hyper-roots (23) depicted with red dots.



(c) 2D rendering of  $J\left(\varphi^{o^2} + \frac{1}{2}\right)$ , s = 0.75

Fig. 10: Visualization of the non divergent set in dynamic space of the function  $\overset{o}{f} = \overset{o}{\varphi}^2 + \frac{1}{2}$  seen from the *s* axis.

set, seen straight from the s axis, is shown in figure 10a. The same viewpoint (slightly magnified), but restricted to a thin layer  $0.74 \le s \le 0.75$ , gives the image shown in 10b. It is finally this last image that resembles the 2D rendering 10c in the hyperimaginary - hyperimaginary plane.

# 8. Conjectures

Differentiability is a necessary condition for a scator function  $\overset{o}{\varphi}(\check{\zeta}) = f_0 + f_x \check{\mathbf{e}}_x + f_y \check{\mathbf{e}}_y$  to be holomorphic. According to a differential quotient criterion, necessary conditions for a function to be differentiable are [Fernández-Guasti, 2018]:

**Theorem 2** [Fernandez-Guasti, 2018]. If a scator function  $\overset{o}{\varphi} : U \subseteq \mathbb{S}^{1+2} \to \mathbb{S}^{1+2}$  of scator variable  $\overset{o}{\zeta}$  in  $U \subseteq \mathbb{S}^{1+2}$ , is differentiable at the point  $\overset{o}{\zeta}_p = z_{0p} + z_{xp} \check{\mathbf{e}}_x + z_{yp} \check{\mathbf{e}}_y$ , then the scalar part of the function  $f_0$ 

and the director parts of the function  $f_x, f_y$ , are real differentiable  $C^1$  functions that satisfy the partial differential equations

$$\frac{\partial f_0}{\partial z_0} = \frac{\partial f_x}{\partial z_x} = \frac{\partial f_y}{\partial z_y},\tag{24a}$$

$$\frac{\partial f_x}{\partial z_0} = -\frac{\partial f_0}{\partial z_x}, \quad \frac{\partial f_y}{\partial z_0} = -\frac{\partial f_0}{\partial z_y}$$
(24b)

$$\frac{\partial f_0}{\partial z_x} \frac{\partial f_0}{\partial z_y} = -\frac{\partial f_x}{\partial z_x} \frac{\partial f_x}{\partial z_y},\tag{24c}$$

evaluated at the point  $\overset{o}{\zeta}_{p}$ .

The partial derivatives involved in (24a) and (24b), resemble the Cauchy-Riemann conditions extended to a real and two hyperimaginary dimensions. The square function does not satisfy conditions (24a), (24b) and (24c). Therefore, the square function is not holomorphic anywhere in the scator set. However, it is scator differentiable for points  $\zeta_p^{o}$  with y = 0, or x = 0 since then all crossed terms are equal to zero. This result is expected because the  $s + x \, \check{\mathbf{e}}_x + 0 \, \check{\mathbf{e}}_y$  or the  $s + 0 \, \check{\mathbf{e}}_x + y \, \check{\mathbf{e}}_y$  planes are identical to the complex plane where the square function is complex holomorphic. The lack of holomorphy according to a differential quotient criterion of the square function prevents us from evaluating its derivatives and their subsequent computation at the critical points. The dynamics in a neighborhood of a periodic point. Thus, the classical fractal classification of points into super-attractive, attractive, indifferent or repelling, depending on the value of its derivative, cannot be readily performed for the quadratic mapping in the  $\mathbb{S}^{1+2}$  scator realm.

In  $S^{1+2}$  it is not possible, according to the differential quotient criterion, to define Fatou or Julia sets in terms of normal meromorphic families, because the scator quadratic function is not meromorphic. Earlier on we mentioned zero and infinity,

Sequences not in the J set either go to  $\overset{o}{\varphi}_{\mathbb{S}^{1+0}} = \frac{1}{2} - \frac{\sqrt{3}}{2} \approx -0.366$ , or diverge. Tested with many points, 25 iterations point 1, .3, .2 goes to -0.366514,  $-4.17295 * 10^{-9}$ , -0.000873619, inter-dyn-sols-1.nb. Convergence is faster in s1+2 point 1, .3, 0 goes to -0.368041, 0.00092023, 0.

conjecture: The Ix-set, provided that nilpotent points are discarded, is the set of parameters where the J set is connected.

Fundamental dichotomy - either connected or infinite separate sets, Cantor dust

The scator set disjointly splits into two sets: the closure of the repelling periodic points and the open set of normal (i.e. stable) points.

Case 1. Nilpotent points

$$r^{4} - 2\sqrt{2}sr^{3} - 4s^{2}r^{2} + 2\sqrt{2}\left(2s^{3} + s^{2}\right)r + \left(4s^{4} + 4s^{3}\right) = 0.$$
(25)

lemma: There is no upper bound b to the scator squared magnitude, such that scators with magnitude larger than b diverge under the quadratic iteration in parameter space.

## 9. Conclusions

Dynamics of fixed points

approaches the fixed point

The action of the other three roots on the  $\varphi_{inv1}^{o}$  fixed point.

inverse orbit can be evaluated with one of the four possible roots.

richness of points contrasted with inverse orbits of fixed points of  $c = -\frac{1}{2}$  in  $\mathbb{C}$ .

The **S** set in  $\overline{\mathbb{E}}^{1+2}_{-}(s; x, y)$  has been defined in parameter space in terms of scators whose magnitude does not tend to infinity under the quadratic iteration. Imaginary scator algebra is a finite dimensional algebra

#### REFERENCES 19

with the peculiarity that the scator product is commutative but does not distribute over addition. It is equipped with an order parameter that in addition to the sum of the squared components it also involves terms with the inverse squared of the scalar component. The quadratic mapping has been justified on two grounds: i) The scator square function is defined by the scator product operation of an element with itself, and ii) If the argument of the square function is multiplied by a scalar, the outcome is equivalent to multiplication by the square of the scalar (Lemma ??). The 3D S set exhibits a rich and intricate boundary not found in other higher dimensional generalizations of the Mandelbrot set.

# Acknowledgements

The scator iteration code has been implemented in the Mandelbulber three dimensional fractal visualization program by the developer's team, in particular, Krzysztof Marczak and Graeme McLaren.

# References

Araki, Y. [2006] "Materializing 3d quasi-fuchsian fractals," Forma 21, 19–27.

- Aron, J. [2009] "The mandelbulb: first 'true' 3d image of famous fractal," New Scientist 204, 54.
- Bedding, S. & Briggs, K. [1995] "Iteration of quaternion maps," International Journal of Bifurcation and Chaos 05, 877–881, doi:10.1142/S0218127495000661, URL http://www.worldscientific.com/doi/abs/ 10.1142/S0218127495000661.
- Blackledge, J. M. [2002] Fractal Geometry: Mathematical Methods, Algorithms, Applications (Woodhead Publishing), ISBN 978-1904275008.
- Bonzini, P. [2010] "To quaternions and back," http://www.fractal.org/mbulb-paolo-bonzini.pdf.
- Cheng, J. & Tan, J. R. [2007] "Generalization of 3d mandelbrot and julia sets," *Journal of Zhejiang University Science A* 8, 134–141, doi:10.1631/jzus.2007.A0134.
- Fernández-Guasti, M. [2016] "Imaginary scators bound set under the iterated quadratic mapping in 1+2 dimensional parameter space," *Int. J. of Bifurcation and Chaos* 26, 1630002, URL https://luz.izt.uam.mx/mfg/arti/16-20/frac-imsca-intro-ijbc-7final.pdf.
- Fernández-Guasti, M. [2018] "Differential quotients in elliptic scator algebra," Math. Meth. App. Sci. 41, 4827–4840, URL https://luz.izt.uam.mx/mfg/arti/16-20/difquot-scaim28-mmas-corr1.pdf.
- Fernández-Guasti, M. [2021a] "Powers of elliptic scator numbers," Axioms 10, 250, doi:10.3390/ axioms10040250, URL https://doi.org/10.3390/axioms10040250.
- Fernández-Guasti, M. [2021b] "Roots of second order polynomials with real coefficients in elliptic scator algebra," J. New Theory 36, 39–48, doi:10.53570/jnt.956340, URL https://doi.org/10.53570/jnt.956340.
- Fernández-Guasti, M. [2022] "Multiplicity of scator roots and the square roots in  $\mathbb{S}^{1+2}$ ," J. New Theory Submitted.
- Fernández-Guasti, M. & Zaldívar, F. [2013] "An elliptic non distributive algebra," Adv. Appl. Clifford Algebras 23, 825–835, URL https://luz.izt.uam.mx/mfg/arti/11-15/imsca-aaca2013.pdf.
- Gomatam, J., Doyle, J., Steves, B. & McFarlane, I. [1995] "Generalization of the mandelbrot set: Quaternionic quadratic maps," *Chaos, Solitons & Fractals* 5, 971–986, doi:10.1016/0960-0779(94)00163-K, URL http://www.sciencedirect.com/science/article/pii/096007799400163K.
- Kantor, I. L. & Solodovnikov, A. S. [1989] *Hypercomplex numbers* (Springer-Verlag), ISBN 978-0-387-96980-0, translated by A. Shenitzer.
- Rama, B. & Mishra, J. [2011] "Generation of 3D fractal images for Mandelbrot set," Proceedings of the 2011 International Conference on Communication, Computing & Security, ICCCS '11 (ACM, New York, NY, USA), ISBN 978-1-4503-0464-1, p. 235–238, doi:10.1145/1947940.1947990, URL http:// doi.acm.org/10.1145/1947940.1947990.
- Sanderson, K. [2009] "2009 gallery: Images of the year," Nature 462, 972–977, doi:10.1038/462972a.
- White, D. & Nylander, P. [2009] "Triplex algebra," http://www.fractalforums.com/theory/triplexalgebra/.